

# US-GUIDED FEMORAL AND SCIATIC NERVE BLOCKS FOR ANALGESIA DURING ENDOVENOUS LASER ABLATION

Prof Saim Yilmaz, MD, Department of Radiology, Akdeniz University School of Medicine, Antalya-Turkey

## Purpose

Endovenous laser ablation may be associated with significant pain when performed under standard local tumescent anesthesia. The purpose of this study was to investigate the value of femoral and sciatic nerve blocks for analgesia during endovenous ablation in patients with lower extremity venous insufficiency.

## Materials/Methods

Between June 2009 and March 2010, US-guided femoral and sciatic nerve blocks were performed to provide analgesia during ELA in 234 legs in 163 patients. The demographic and clinical aspects of the patients are given in table 1. For femoral block, the patient was put supine. After the groin was disinfected, the femoral vessels were visualized in transverse section. The femoral nerve was identified in the hyperechoic triangle formed by the lateral border of the common femoral artery, fascia iliaca and iliopsoas muscle. Then, a 3-4ml 2% lidocaine or bupivacaine solution diluted to 10ml with saline was injected into this triangle under US guidance (Figure 1). For sciatic block, the patient was put prone. The posterior aspect of the knee and lower thigh were disinfected. The popliteal vessels were visualized in the popliteal fossa in transverse section. The sciatic nerve was first identified posterolateral to the popliteal artery at the knee, and then scanned upto the midthigh level. Then, the same amount of lidocaine or bupivacaine solution was injected around the nerve under US guidance (Figure 2).

After the blocks, ELA procedure was performed using the standard tumescent anesthesia. In 234 legs, 255 refluxing veins were ablated. Femoral block was used for the ELA of 191 GSVs and 10 perforating veins, and sciatic block was used for the ELA of 46 SSVs and 8 perforating veins. Following the ELA, concomitant foam sclerotherapy was also performed in 220 of 234 legs. After the procedures, the patient was put on compression stockings and the motor function was evaluated. In the presence of persistent motor block, a short-stretch elastic bandage was applied around the knee to prevent sudden flexion and the patient was accompanied by another person during the walking until the motor function was completely recovered.

After the procedures, a visual analogue scale (1–10) was used for pain assessment (Figure 3).

## Results

After the blocks, pain scores were 0 in 99 legs, 1 in 61, 2 in 44, 3 in 23 and 4 in 7 legs. After femoral nerve block, pain scores were lower in the left leg compared to the right (Table 2), and this difference was statistically significant ( $p=0,000$ ). After sciatic block, pain scores were lower compared to the femoral block (table 2), and this difference was also statistically significant ( $p=0,000$ ). After femoral block, mild to moderate motor block occurred in 18 legs (14 in the left, 4 in the right). Venous spasm was observed in only 2 legs (in the right) during the catheterization.

**Table 1: Demographic and clinical aspects of the study population.**

Number of patients	163, 234 legs (bilateral in 71 patients)
Age	23-66 years
Sex	127 female, 36 male
Incompetent veins	255 (191 GSV, 46 SSV, 18 Perforating veins)
Clinical classification (CEAP) in 234 legs	C1 (n=10) C2 (n=141) C3 (n=24) C4a (n=20) C4b (n=16) C5 (n=12) C6 (n=11)

**Table 2: Technical details and results of US-guided nerve blocks.**

Extremity	111 right, 123 left
Nerve block	Femoral in 192, Sciatic in 50 (both in 8 legs)
Total pain scores in the right and left legs	137 in the right leg (0 in 39, 1 in 28, 2 in 27, 3 in 13, 4 in 4), <b>Mean: 1.23±0.11</b> 109 in the left leg (0 in 60, 1 in 33, 2 in 17, 3 in 10, 4 in 3) <b>Mean: 0.89±0.08 (P=0,000)</b>
Total pain scores in femoral and sciatic blocks	200 in femoral (0 in 83, 1 in 50, 2 in 33, 3 in 20, 4 in 6), <b>Mean: 1.04±0.18</b> 42 in sciatic (0 in 26, 1 in 11, 2 in 9, 3 in 3, 4 in 1) <b>Mean: 0.84±0.15 (P=0,000)</b>




Figure 1. US-guided femoral nerve block: A. For femoral block, the patient is put supine and the transducer is held parallel to the inguinal ligament and femoral vessels are visualized in transverse section. B. Femoral nerve (FN) is within the echogenic triangular area lateral to the common femoral artery (CFA) and vein (CFV).




Figure 2. US-guided sciatic nerve block: A. For sciatic block, the patient is put prone and the transducer is held proximal to the the popliteal fossa and popliteal vessels are visualized in transverse section. B. Sciatic nerve (SN) is seen as a large, round and echogenic structure posterolateral to the popliteal artery (PA) and vein (PV).




Figure 3: Visual analogue pain scale

## Conclusion

ELA is a well-accepted alternative to surgery in the treatment of saphenous and perforating venous insufficiency. The procedure is normally performed using only local tumescent anesthesia. The aim of tumescent anesthesia is not only to eliminate pain during ELA, but also to protect the surrounding tissues from laser and to empty the vein, bringing the vein wall in close contact with the laser fiber (1-3). However, multiple needle punctures and injection of the local anesthetic solution along the veins such as GSV and SSV may induce considerable pain during the tumescent anesthesia. Although the intensity of the pain is "tolerable" for most patients, it may be quite a "bad experience" for the others. Pain may be particularly intense if the patient develops venous spasm during the catheterization. Additional pain may also be created if foam sclerotherapy is performed concomitantly following ELA (4).

A number of methods are currently used to decrease pain during the ELA. Some physicians, perform the procedure under spinal or even general anesthesia. Although the patient has "no pain" with these methods, they are generally not recommended because: 1. Delayed mobilization may increase the risk of deep venous thrombosis. 2. Deep anesthesia may increase the risk of saphenous or sural nerve injury during ELA (5,6). 3. The cost is increased since the procedure requires dedicated staff and a hospital stay. Other physicians use oral or parenteral pain killers to cope with the pain problem. Our personal experience does not favor this method, because: 1. Standard NSAIDs are generally sufficient for postoperative pain, but they are not very effective for pain during the procedure. 2. Narcotic analgesics are more effective, but may cause hypotension and other hemodynamic problems and thus, interfere with the mobility of the patient after the procedure.

Alternatively, US-guided femoral or sciatic nerve blocks may be used for analgesia during ELA (7). In our experience, this method was very effective in the elimination of pain during the ELA procedure. In our study, we did not compare the pain scores of patients in whom we performed US-guided nerve blocks versus those in whom we did not. However, very low pain scores of the patients suggest that US-guided femoral and sciatic nerve blocks are highly effective in the elimination of pain during the ELA. In our patients, our aim was to provide analgesia while preserving the motor function, keeping the patient as active as possible after the treatment. To do this, we diluted the local anesthetic with saline and chose a relatively short-acting local anesthetic (lidocaine in most patients). Despite that, transient mild to moderate motor block occurred after femoral nerve blockage in 18 legs (7.6%). In these, we applied a bandage around the knee to prevent unintentional flexion and the patient was accompanied by another person during the walking until the motor function was completely recovered. In our study, the femoral block was more effective in the left leg compared to the right (pain scores were significantly lower and the frequency of motor block was higher). Although we don't have a definitive explanation for this difference, it may be that in right-handed individuals (the majority of the general population), peripheral nerves on the left side may be more susceptible to local anesthetics.

In our experience, another advantage of femoral and sciatic nerve blocks was to decrease the frequency of venous spasm. In the patients we performed ELA solely with tumescent anesthesia, we encountered venous spasm relatively frequently, particularly when the vein was thin and when nonhydrophilic guidewires and catheters were used. In the present study, although we used the same materials, we saw venous spasm in only two patients. In the remaining patients, there was no venous spasm during the ELA, and we observed a slight increase in the diameter of the refluxing veins after the nerve block, probably due to the sympathetic blockage. In our experience, this venous distension made the puncture and catheterization easier, and together with the absence of venous spasm, markedly facilitated the ELA procedure.

Finally, femoral and sciatic nerve blocks are easy to perform for phlebologists who are familiar with US-guided interventions. The procedures do not require any additional equipment and staff; all the medications, materials and US machine are already used during the ELA. Thus, there is no additional cost. The amount of local anesthetic is far lower than the toxic limit. Thus, the procedure is extremely safe and can be performed in the office setting.

In conclusion, US-guided femoral and sciatic nerve blocks may provide considerable reduction in pain and venous spasm during endovenous laser ablation. They may make the procedure more comfortable for the patient and easier for the operator.

## References

- Min RJ, Khilnani NM. Endovenous laser ablation of varicose veins. J Cardiovasc Surg. 2005; 46: 395-405.
- Yilmaz S, Ceken K, Alparslan A, Sindel T, Lüleci E. Endovenous laser ablation for saphenous vein insufficiency: immediate and short-term results of our first 60 procedures. Diagn Interv Radiol. 2007; 13:156-163.
- van den Bos R, Arends L, Kockaert M, Neumann M, Nijsten T. Endovenous therapies of lower extremity varicosities: a meta-analysis. J Vasc Surg. 2009; 49: 230-239.
- Coleridge Smith P. Sclerotherapy and foam sclerotherapy for varicose veins. Phlebology. 2009; 24: 260-269.
- Marsh P, Price BA, Holdstock J, Harrison C, Whiteley MS. Deep Vein Thrombosis (DVT) after Venous Thermoablation Techniques: Rates of Endovenous Heat-induced Thrombosis (EHIT) and Classical DVT after Radiofrequency and Endovenous Laser Ablation in a Single Centre. Eur J Vasc Endovasc Surg. 2010 Jul 22. E pub.
- Van Den Bos RR, Neumann M, De Roos KP, Nijsten T. Endovenous laser ablation-induced complications: review of the literature and new cases. Dermatol Surg. 2009 35: 1206-1214.
- Fingerman M, Benonis JG, Martin G. A practical guide to commonly performed ultrasound-guided peripheral-nerve blocks. Curr Opin Anaesthesiol. 2009; 22: 600-607.